

## Contents

<b>Introduction</b>	5
<b>Chapter 1.</b>	
<b>Physical concepts of Mössbauer spectroscopy</b>	11
<i>1.1. Recoilless nuclear resonance emission and absorption</i>	11
<i>1.2. Temperature shift of the Mössbauer spectrum</i>	19
<i>1.3. Isomeric Shift of the Mössbauer spectrum</i>	19
<i>1.4. Quadrupole splitting</i>	21
<i>1.5. Emission Mössbauer spectroscopy of the after-effects of nuclear transformations in solids</i>	25
<i>1.6. One-electron exchange between neutral and ionized acceptor iron centers in GaAs and GaP</i>	27
<i>1.7. Observation of charge exchange by copper impurity atoms in AgCl</i>	38
<i>1.8. Tin chalcogenides</i>	41
<i>1.9. Compounds of tin with elements of the fifth group</i>	45
<i>1.10. Tin impurity atoms in silicon, germanium, and silicon-germanium solid</i>	53
<b>Chapter 2.</b>	
<b>Identification of neutral and ionized tin donor states in PbS and PbSe</b>	64
<i>2.1. Charge exchange of tin atoms in <math>Pb_{1-x}Sn_xS</math> solid solutions</i>	64
<i>2.2. Charge exchange of tin atoms in <math>Pb_{1-x}Sn_xSe</math> solid solutions</i>	67
<i>2.3. Local symmetry and electronic structure of atoms in the <math>Pb_{1-x-y}Sn_xNa_ySe</math> lattices in the gapless state</i>	70
<i>2.4. Two-electron tin centers with negative correlation energy in lead chalcogenides. Determination of the Hubbard energy</i>	78
<i>2.5. Hubbard energy of two-electron tin centers in <math>PbS_{1-z}Te_z</math> solid solutions</i>	89

2.6. Local symmetry and electronic structure of tin atoms in $(\text{Pb}_{1-x}\text{Sn}_x)_{1-z}\text{In}_z\text{Te}$ lattices	92
2.7. Statistics of electrons in PbS with U centers	96
2.8. Energy parameters of two-electron tin centers in PbSe	104
2.9. Detection of singly ionized state of two-electron tin centers with negative correlation energy in $\text{Pb}_{1-x}\text{Sn}_x\text{S}$ alloys	112
2.10. Two-electron centers with negative correlation energy $\text{Pb}_{1-x}\text{Sn}_x\text{Se}$ solid solutions	117
<b>Chapter 3.</b>	
<b>Two-electron tin centers formed in lead chalcogenides as a result of nuclear transmutations</b>	123
3.1. Position of antimony impurity atoms in a PbTe lattice, determined by emission Mössbauer spectroscopy	123
3.2. Antistructural defects in PbTe-type semiconductors	127
3.3. Determining the position of antimony impurity atoms in PbS by $^{119}\text{Sb}(^{119m}\text{Sn})$ emission Mössbauer spectroscopy	131
3.4. Two-electron tin centers formed in lead chalcogenides as a result of nuclear transmutations	135
3.5. States of antimony and tin atoms in lead chalcogenides	148
3.6. Two-electron tin centers with a negative correlation energy in PbS	156
<b>Chapter 4.</b>	
<b>Identification of tin states in chalcogenide glasses</b>	172
4.1. The influence of the crystal- glass transition on the state of impurity tin atoms in chalcogenide semiconductors	172
4.2. Tin impurity atom states in modified $\text{As}_2\text{Se}_3$ films	181
4.3. Properties and structure of $(\text{As}_2\text{Se}_3)_{1-z}(\text{SnSe})_{z-x}(\text{GeSe})_x$ and $(\text{As}_2\text{Se}_3)_{1-z}(\text{SnSe}_2)_{z-x}(\text{GeSe}_2)_x$ glasses	186
4.4. Local structure of Ge – S, Ge – Se, and Ge – Te vitreous alloys	197

4.5. Structure and physicochemical properties of glasses in the $(As_2Se_3)_{1-z}(SnSe_2)_{z-x}(Te_2Se)_x$ and $(As_2Se_3)_{1-z}(SnSe)_{z-x}(Te_2Se)_x$ systems	202
<b>Chapter 5.</b>	
<b>Two-electron tin centers formed in chalcogenide glasses as a result of nuclear transmutations</b>	214
5.1. Two-electron tin centers formed in glasses $As_2S_3$ , $As_2Se_3$ , and $As_2Te_3$ as a result of nuclear transformations	214
5.2. Impurity centers of tin in glassy arsenic chalcogenides	224
5.3. The state of tin impurity atoms in vitreous germanium chalcogenides	233
<b>Chapter 6.</b>	
<b>Electron exchange between impurity centers of tin in lead chalcogenides and chalcogenide glasses</b>	247
6.1. Identification of neutral and ionized tin donor states and the observation of two-electron exchange between tin centers in solid solutions based on $PbS$ and $PbSe$	247
6.2. Mechanism of two-electron exchange between neutral and ionized centres of tin in $PbS_{1-x}Se_x$ solid solutions	257
6.3. Investigation of two-electron exchange between neutral and ionized impurity centers	266
6.4. Unconventional tin atom states in mixed silver and tin chalcogenides having $NaCl$ structure	274
6.5. Temperature dependence of the frequency of two-electron exchange between impurity negative- $U$ tin centers in lead sulfide	279
6.6. Electron exchange between impurity centers of tin in lead chalcogenides	283
6.7. Electron exchange between tin impurity $U^-$ centers in $PbS_zSe_{1-z}$ alloys	294
6.8. The two-electron exchange between the $U^-$ tin centers in crystal and Glass-like chalcogenide semiconductors	307

<b>Chapter 7.</b>	
<b>The electrical activity of germanium impurities in lead chalcogenides</b>	312
<i>7.1. The electrical activity of isoelectronic germanium impurities in lead chalcogenides</i>	312
<i>7.2. Arrangement of arsenic atoms in the PbTe lattice</i>	317
<i>7.3. Observation of Bose condensation in <math>(\text{Pb}_{0.4}\text{Sn}_{0.6})_{0.86}\text{In}_{0.14}\text{Te}</math> semiconductor solid solutions using Mössbauer spectroscopy</i>	321
<i>7.4. Two-electron germanium centers with a negative correlation energy in lead chalcogenides</i>	327
<i>7.5. Mössbauer <math>\text{U}^-</math> centers as tools for studying the Bose condensation in semiconductors</i>	339
<b>Conclusions</b>	351
<b>Appendix</b>	
<b>Determination of the composition of chalcogenide semiconductors X-Ray Fluorescence Analysis</b>	352
<i>A.1. Determination of the composition of binary chalcogenide glasses by X-Ray fluorescence analysis</i>	352
<i>A.2. Determination of the composition of multicomponent chalcogenide semiconductors by x-ray fluorescence analysis</i>	360
<i>A.3. X-ray fluorescence analysis of <math>\text{Ge}_{1-x}\text{Se}_x</math>, <math>\text{As}_{1-x}\text{Se}_x</math>, and <math>\text{Ge}_{1-x-y}\text{As}_y\text{Se}_x</math> glasses using electronic excitation</i>	372
<b>References</b>	383